

(H, G) -coincidence theorems for manifolds and a topological Tverberg type theorem for any natural number r

Denise de Mattos*, Edivaldo L. dos Santos[†] and Taciana O. Souza[‡]

Abstract

Let X be a paracompact space, let G be a finite group acting freely on X and let H a cyclic subgroup of G of prime order p . Let $f : X \rightarrow M$ be a continuous map where M is a connected m -manifold (orientable if $p > 2$) and $f^*(V_k) = 0$, for $k \geq 1$, where V_k are the Wu classes of M . Suppose that $\text{ind } X \geq n > (|G| - r)m$, where $r = \frac{|G|}{p}$. In this work, we estimate the cohomological dimension of the set $A(f, H, G)$ of (H, G) -coincidence points of f . Also, we estimate the index of a (H, G) -coincidence set in the case that H is a p -torus subgroup of a particular group G and as application we prove a topological Tverberg type theorem for any natural number r . Such result is a weak version of the famous topological Tverberg conjecture, which was proved recently, fail for all r that are not prime powers. Moreover, we obtain a generalized Van Kampen-Flores type theorem for any natural number r .

2010 *Mathematics Subject Classification*: Primary 55M20, 52A35; Secondary 55M35, 55S35.

1 Introduction

Let G be a finite group which acts freely on a space X and let $f : X \rightarrow Y$ be a continuous map from X into another space Y . If H is a subgroup of G , then H acts on the right on each orbit Gx of G as follows: if $y \in Gx$ and $y = gx$, with $g \in G$, then $h \cdot y = gh^{-1}x$. A point $x \in X$ is said to be a (H, G) -coincidence point of f (as introduced by Gonçalves, Pergher and Jaworowski in [8]) if f sends every orbit of the action of H on the G -orbit of x to a single point. Of course, if H is the trivial subgroup, then every point of X is a (H, G) -coincidence. If $H = G$, this is the usual definition of G -coincidence, that is, $f(x) = f(gx)$, for all $g \in G$. If $G = \mathbb{Z}_p$ with p prime, then

*The author was supported in part by FAPESP of Brazil Grant numbers 12/24454-8 and 2013/24845-0

[†]The author was supported in part by FAPESP of Brazil Grant numbers 12/24454-8

[‡]The author was supported by CAPES and CNPq

a nontrivial (H, G) -coincidence point is a G -coincidence point. Let us denote by $A(f, H, G)$ the set of all (H, G) -coincidence points. Borsuk-Ulam type theorems consists in estimating the cohomological dimension of the set $A(f, H, G)$. Two main directions considered of this problem are either when the target space Y is a manifold or Y is a CW complex. In the first direction are the papers of Borsuk [4] (the classical theorem of Borsuk-Ulam, for $H = G = \mathbb{Z}_2$, $X = S^n$ and $Y = \mathbb{R}^n$), Conner and Floyd [5] (for $H = G = \mathbb{Z}_2$ and Y a n -manifold), Munkholm [13] (for $H = G = \mathbb{Z}_p$, $X = S^n$ and $Y = \mathbb{R}^m$), Nakaoka [14] (for $H = G = \mathbb{Z}_p$, X under certain (co)homological conditions and Y a m -manifold) and the following more general version proved by Volovikov [17] using the index of a free $\mathbb{Z}_p$ -space X ($\text{ind } X$, see Definition 2.2):

Theorem A.[17, Theorem 1.2] *Let X be a paracompact free $\mathbb{Z}_p$ -space of $\text{ind } X \geq n$, and $f : X \rightarrow M$ a continuous mapping of X into an m -dimensional connected manifold M (orientable if $p > 2$). Assume that:*

- (1) $f^*(V_i) = 0$ for $i \geq 1$, where the V_i are the Wu classes of M ; and
- (2) $n > m(p - 1)$.

Then the $\text{ind } A(f) \geq n - m(p - 1) > 0$.

In the second direction are the papers of Izydorek and Jaworowski [10] (for $H = G = \mathbb{Z}_p$, $X = S^n$ and Y a CW-complex), Gonçalves and Pergher [7] (for $H = G = \mathbb{Z}_p$, $X = S^n$ and Y a CW-complex) and for proper nontrivial subgroup H of G , Gonçalves, Jaworowski and Pergher [8] (for $H = \mathbb{Z}_p$ subgroup of a finite group G , X an homotopy sphere and Y a CW-complex) and Gonçalves, Jaworowski, Pergher and Volovikov [9](for $H = \mathbb{Z}_p$ subgroup of a finite group G , X under certain (co)homological assumptions and Y a CW-complex).

In this work, considering the target space $Y = M$ a manifold and H a proper nontrivial subgroup of G , we prove the following formulation of the Borsuk-Ulam theorem for manifolds in terms of (H, G) -coincidence.

Theorem 1.1. *Let X be a paracompact space of $\text{ind } X \geq n$ and let G be a finite group acting freely on X and H a cyclic subgroup of G of prime order p . Let $f : X \rightarrow M$ be a continuous map where M is a connected m -manifold (orientable if $p > 2$) and $f^*(V_k) = 0$, for all $k \geq 1$, where V_k are the Wu classes of M . Suppose that $\text{ind } X \geq n > (|G| - r)m$ where $r = \frac{|G|}{p}$, then $\text{ind } A(f, H, G) \geq n - (|G| - r)m$. Consequently,*

$$\text{cohom.dim } A(f, H, G) \geq n - (|G| - r)m > 0.$$

Let us observe that if $H = G = \mathbb{Z}_p$, we have $(|G| - r)m = (p - 1)m$ and therefore Theorem 1.1 generalizes Theorem A. obtained by Volovikov in [17] in terms of (H, G) -coincidence. For the case $n = (|G| - r)m$, p an odd prime, if we consider X a $\text{mod } p$ homology n -sphere in the Theorem 1.1(in this case, the continuous map f can be arbitrary), it follows from [14, Theorem 8] that

$A(f, H, G) \neq \emptyset$. Also, Theorem 1.1 is a version for manifolds of the main result due to Gonçalves, Jaworowski and Pergher in [8].

Also, we prove the following nonsymmetric theorem for (H, G) -coincidences which is a version for manifolds of the main theorem in [11].

Theorem 1.2. *Let X be a compact Hausdorff space, let G be a finite group acting freely on S^n and let H be a cyclic subgroup of G of order prime p . Let $\varphi : X \rightarrow S^n$ be an essential map¹ and let $f : X \rightarrow M$ be a continuous map where M is a connected m -manifold (orientable if $p > 2$) and $f^*(V_k) = 0$, for all $k \geq 1$, where V_k are the Wu classes of M . Suppose that $n > (|G| - r)m$, then*

$$\text{cohom.dim } A_\varphi(f, H, G) \geq n - (|G| - r)m,$$

where $r = \frac{|G|}{p}$ and $A_\varphi(f, H, G)$ denotes the (H, G) -coincidence points of f relative to an essential map $\varphi : X \rightarrow S^n$.

In Section 5, we give similar estimate in the case that H is a p -torus subgroup of a particular group G and as application, we prove a topological Tverberg type theorem for any natural number, which is a weak version of the famous topological Tverberg conjecture. Moreover, we obtain a generalized Van Kampen-Flores type theorem for any integer r .

2 Preliminaries

We start by introducing some definitions as follows.

2.1 The $\mathbb{Z}_p$ -index

We suppose that the cyclic group $\mathbb{Z}_p$ acts freely on a paracompact Hausdorff space X , where p is a prime number and we denote by $[X]^*$ the space of orbits of X by the action of $\mathbb{Z}_p$. Then, $X \rightarrow [X]^*$ is a principal $\mathbb{Z}_p$ -bundle and we can consider a classifying map $c : [X]^* \rightarrow B\mathbb{Z}_p$.

Remark 2.1. It is well known that if $\hat{c}$ is another classifying map for the principal $\mathbb{Z}_p$ -bundle $X \rightarrow X^*$, then there is a homotopy between c and $\hat{c}$.

Definition 2.2. *We say that the $\mathbb{Z}_p$ -index of X is greater than or equal to l if the homomorphism*

$$c^* : H^l(B\mathbb{Z}_p; \mathbb{Z}_p) \rightarrow H^l([X]^*; \mathbb{Z}_p)$$

is nontrivial. We say that the $\mathbb{Z}_p$ -index of X is equal to l if it is greater or equal than l and, furthermore, $c^* : H^i(B\mathbb{Z}_p; \mathbb{Z}_p) \rightarrow H^i([X]^*; \mathbb{Z}_p)$ is zero, for all $i \geq l + 1$.

We denote the $\mathbb{Z}_p$ -index of X by $\text{ind } X$.

¹A map $\varphi : X \rightarrow S^n$ is said to be an essential map if φ induces nonzero homomorphism $\varphi^* : H^n(S^n; \mathbb{Z}_p) \rightarrow H^n(X; \mathbb{Z}_p)$.

2.2 Wu classes

The *total Wu class* of a manifold M is defined as the formal sum

$$v(M) = 1 + v_1(M) + v_2(M) + \cdots + v_k(M) + \cdots$$

where $v_k(M)$ is the k -th Wu class of M , $k = 1, 2, \dots$ (see [12]). Let $p > 2$ be a prime, using the *total reduced power*

$$P = P^0 + P^1 + P^2 + \cdots + P^k + \cdots$$

and the equation

$$\langle v_k(M) \smile x, [M] \rangle = \langle P^k(x), [M] \rangle$$

we obtain the formula

$$\langle v(M) \smile u, [M] \rangle = \langle P(u), [M] \rangle$$

for all $u \in H_c^*(M; \mathbb{Z}_p)$. For $p = 2$ we have a similar formula

$$\langle v(M) \smile u, [M] \rangle = \langle Sq(u), [M] \rangle$$

for all $u \in H_c^*(M; \mathbb{Z}_2)$, where

$$Sq = Sq^0 + Sq^1 + Sq^2 + \cdots + Sq^k + \cdots$$

is the *total Steenrod square*.

3 The Wu class for product of manifolds

Let W and M be manifolds with dimensions w and m respectively, $w \geq m$, both orientables if $p > 2$. In the next lemma, we obtain a characterization of the Wu class of the product of W and M .

Lemma 3.1. *Let W and M be a connected manifolds, both orientables if $p > 2$. Then, the total Wu class of $W \times M$, is given by:*

$$v(W) \otimes v(M) \tag{3.1}$$

where $v(W)$ and $v(M)$ are the total Wu classes of W and M respectively.

Proof. Let $p > 2$ be a prime number. Let $z = w \otimes u$ an element of $H_c^*(W \times M; \mathbb{Z}_p)$ then

$$\begin{aligned} \langle v(W) \otimes v(M) \smile z, [W \times M] \rangle &= \langle v(W) \smile w \otimes v(M) \smile u, [W \times M] \rangle \\ &= \langle P(w) \otimes P(u), [W \times M] \rangle \\ &= \langle P(w \otimes u), [W \times M] \rangle \\ &= \langle P(z), [W \times M] \rangle \\ &= \langle v(W \times M) \smile z, [W \times M] \rangle \end{aligned}$$

Therefore by uniqueness of the Wu class we conclude that the total Wu class of $W \times M$ is given by $v(W \times M) = v(W) \otimes v(M)$. By a similar argument the total Wu classes are obtained for $p = 2$, in this case are used the total Steenrod square. ■

4 (H, G) -Coincidence theorems for manifolds

Now, we denote by $a_1, \dots, a_r$ a set of representatives of the left lateral classes of G/H . We define, for each $i = 1, \dots, r$, $g_i : X \rightarrow X$ by $g_i(x) = a_i x$. Consider the map $F : X \rightarrow M^r$ defined by

$$F = (f_1 \times \dots \times f_r) \circ D,$$

where $D : X \rightarrow X^r$ is the diagonal map and $f_i = f \circ g_i$. We prove the following

Lemma 4.1. *If $f^*(v_k(M)) = 0$, for all $k \geq 1$, where $v_k(M)$ are the Wu classes of M , then $F^*(v_k(M^r)) = 0$, for all $k \geq 1$, where $v_k(M^r)$ are the Wu classes of M^r .*

Proof. It suffices to show that $(f_1 \times \dots \times f_r)^*(v_k(M^r)) = 0$, for $k \geq 1$. If $r = 1$, then $F = f_1$ and $f_1^*(v_k(M)) = g_1^* \circ f^*(v_k(M)) = 0$.

Let us denote by

$$p_1 : M^{r-1} \times M \rightarrow M^{r-1}, p_2 : M^{r-1} \times M \rightarrow M$$

$$q_1 : X^{r-1} \times X \rightarrow X^{r-1}, q_2 : X^{r-1} \times X \rightarrow X$$

the natural projections. If $r \geq 2$, we have

$$(f_1 \times \dots \times f_{r-1}) \circ q_1 = p_1 \circ (f_1 \times \dots \times f_r)$$

$$f_r \circ q_2 = p_2 \circ (f_1 \times \dots \times f_r).$$

Since, by Lemma 3.1, $v_k(M^{r-1} \times M) = \sum_{s=0}^k v_s(M^{r-1}) \times v_{k-s}(M)$ and assuming inductively that $(f_1 \times \dots \times f_{r-1})^*(v_s(M^{r-1})) = 0$, for $s \geq 1$, we conclude that

$$\begin{aligned} & (f_1 \times \dots \times f_r)^*(v_k(M^{r-1} \times M)) = \\ &= (f_1 \times \dots \times f_r)^* \left(\sum_{s=0}^k v_s(M^{r-1}) \times v_{k-s}(M) \right) \\ &= \sum_{s=0}^k (f_1 \times \dots \times f_r)^*(p_1^*(v_s(M^{r-1}))) \smile (f_1 \times \dots \times f_r)^*(p_2^*(v_{k-s}(M))) \\ &= \sum_{s=0}^k q_1^* \circ (f_1 \times \dots \times f_{r-1})^*(v_s(M^{r-1})) \smile q_2^* \circ g_r^* \circ f^*(v_{k-s}(M)) \\ &= 0. \end{aligned} \quad \blacksquare$$

4.1 Proof of Theorem 1.1 and its consequences

In this section, we present the proofs of Theorems 1.1 and its consequences, as follows.

Proof of Theorem 1.1. We consider the map F defined previously. We have

$$A(f, H, G) \supset A_F = \{x \in X : F(x) = F(hx), \forall h \in H\}.$$

In fact, let x be a point in the set A_F , then

$$(f(a_1x), \dots, f(a_rx)) = (f(a_1hx), \dots, f(a_rhx)),$$

for all $h \in H$. Thus, $f(a_ix) = f(a_ihx)$, for all $h \in H$ and $i = 1, \dots, r$. According to the definition of the action of H on the orbit Gx , $h^{-1} \cdot a_ix := a_i(h^{-1})^{-1}x = a_ihx \in a_iHx$, for $i = 1, \dots, r$. Thus, f collapses each orbit a_iHx determined by the action of H on a_ix , for $i = 1, \dots, r$, therefore $x \in A(f, H, G)$.

Now we observe that $H \cong \mathbb{Z}_p$ acts freely on X by restriction and by hypothesis $\text{ind } X \geq n > n - (p-1)rm$. By Lemma 4.1, $F^*(v_k) = 0$, for all $k \geq 1$, where v_k are the Wu classes of M^r . Thus, according to [17, Theorem 1.2]

$$\text{ind } A_F \geq n - (p-1)rm = n - (|G| - r)m.$$

Let us consider the inclusion $i : A_F \rightarrow A(f, H, G)$, which is an equivariant map, and so it induces $\bar{i} : [A_F]^* \rightarrow [A(f, H, G)]^*$ a map between the orbit spaces. Therefore, if $c : [A(f, H, G)]^* \rightarrow B\mathbb{Z}_p$ is any classifying map, we have that $c \circ \bar{i} : [A_F]^* \rightarrow B\mathbb{Z}_p$ is a classifying map. Thus,

$$\text{ind } A(f, H, G) \geq \text{ind } A_F \geq n - (|G| - r)m,$$

■

Corollary 4.2. *Let X be a paracompact space and let G be a finite group acting freely on X . Let M be a orientable m -manifold, and p a prime number that divide $|G|$. Suppose that $\text{ind } X \geq n > (|G| - r)m$, where $r = \frac{|G|}{p}$. Then, for a continuous map $f : X \rightarrow M$ such that $f^*(V_k) = 0$, for all $k \geq 1$, where V_k are the Wu classes of M , there exists a non-trivial subgroup H of G , such that*

$$\text{cohom.dim } A(f, H, G) \geq n - (|G| - r)m.$$

Proof. Let p be a prime number such that divide $|G|$. By Cauchy Theorem, there is a cyclic of order p subgroup H of G . Then, we apply Theorem 1.1.

■

Remark 4.3. Let us observe that, if $f^* : H^i(M; \mathbb{Z}_p) \rightarrow H^i(X; \mathbb{Z}_p)$ is trivial, for $i \geq 1$, and p is the smallest prime number dividing $|G|$, then $r = \frac{|G|}{p} \geq \frac{|G|}{q}$, where q can be any other prime number dividing $|G|$. Thus, $n > (|G| - \frac{|G|}{q})m$, therefore for each prime number q dividing $|G|$, there exists a cyclic subgroup of order q , H_q of G such that $\text{ind } A(f, H_q, G) \geq n - (|G| - r)m$.

The following theorem is a version for manifolds of the main result in [8]

Theorem 4.4. *Let G be a finite group which acts freely on n -sphere S^n and let H be a cyclic subgroup of G of prime order p . Let $f : S^n \rightarrow M$ be a continuous map where M be a m -manifold (orientable if $p > 2$). If $n > (|G| - r)m$ where $r = \frac{|G|}{p}$, then*

$$\text{cohom.dim}(A(f, H, G)) \geq n - (|G| - r)m.$$

Proof. Since $n > (|G| - r)m \geq m$, $f^*(V_k) = 0$, for all $k \geq 1$. Moreover, $\text{ind } S^n = n$ and thus we apply the Theorem 1.1. $\blacksquare$

4.2 Proof of Theorem 1.2

Now, let us consider X a compact Hausdorff space and an essential map $\varphi : X \rightarrow S^n$. Suppose G be a finite group of order s which acts freely on S^n and H be a subgroup of order p of G . Let $G = \{g_1, \dots, g_s\}$ be a fixed enumeration of elements of G , where g_1 is the identity of G . A nonempty space X_φ can be associated with the essential map $\varphi : X \rightarrow S^n$ as follows:

$$X_\varphi = \{(x_1, \dots, x_s) \in X^s : g_i \varphi(x_1) = \varphi(x_i), i = 1, \dots, s\},$$

where X^s denotes the s -fold cartesian product of X . The set X_φ is a closed subset of X^s and so it is compact. We define a G -action on X_φ as follows: for each $g_i \in G$ and for each $(x_1, \dots, x_s) \in X_\varphi$,

$$g_i(x_1, \dots, x_s) = (x_{\sigma_{g_i}(1)}, \dots, x_{\sigma_{g_i}(s)}),$$

where the permutation σ_{g_i} , is defined by $\sigma_{g_i}(k) = j$, $g_k g_i = g_j$. We observe that if $x = (x_1, \dots, x_s) \in X_\varphi$ then $x_i \neq x_j$, for any $i \neq j$ and therefore G acts freely on X_φ .

Let us consider a continuous map $f : X \rightarrow M$, where M is a topological space and $\tilde{f} : X_\varphi \rightarrow M$ given by $\tilde{f}(x_1, \dots, x_s) = f(x_1)$,

Definition 4.5. *The set $A_\varphi(f, H, G)$ of (H, G) -coincidence points of f relative to φ is defined by*

$$A_\varphi(f, H, G) = A(\tilde{f}, H, G).$$

Proof of Theorem 1.2. Let $\tilde{f} : X_\varphi \rightarrow M$ given by $\tilde{f}(x_1, \dots, x_s) = f(x_1)$, that is, $\tilde{f} = f \circ \pi_1$, where π_1 is the natural projection on the 1-th coordinate. By hypothesis, $f^*(V_k) = 0$, for all $k \geq 1$, where V_k are the Wu classes of M , then we have $\tilde{f}^*(V_k) = 0$, for all $k \geq 1$. Moreover, the $\mathbb{Z}_p$ -index of X_φ is equal to n by [11] Theorem 3.1. In this way, X_φ and $\tilde{f}$ satisfy the hypothesis of Theorem 1.1 which implies that the $\mathbb{Z}_p$ -index of the set $A(\tilde{f}, H, G)$ is greater than or equal to $n - (|G| - r)m$. By definition, $A_\varphi(f, H, G) = A(\tilde{f}, H, G)$, and then

$$\text{cohom.dim } A_\varphi(f, H, G) \geq n - (|G| - r)m.$$

$\blacksquare$

By a similar argument to that used in the proof of Corollary 4.2 we have the following corollary of Theorem 1.2

Corollary 4.6. *Let X be a compact Hausdorff space and let G be a finite group acting freely on S^n . Let M be a orientable m -manifold and p a prime number dividing $|G|$. Suppose that $n > (|G| - r)m$, where $r = \frac{|G|}{p}$. Then, for a continuous map $f : X \rightarrow M$, with $f^*(V_k) = 0$, for all $k \geq 1$, where V_k are the Wu classes of M , there exists a non-trivial subgroup H of G , such that*

$$\text{cohom.dim } A_\varphi(f, H, G) \geq n - (|G| - r)m.$$

5 Topological Tverberg type theorem

The history of Tverberg theorem begins with a Birch's paper (see [2]) which contained the following conjecture

“Any $(r - 1)(d + 1) + 1$ points in $\mathbb{R}^d$ can be partitioned in N subsets whose convex hulls have a common point ”.

The Birch's conjecture was proved by Helge Tverberg (see [16]) and since then is known as Tverberg theorem.

We note that the convex hull of $l + 1$ points in $\mathbb{R}^d$ is the image of the linear map $\Delta_l \rightarrow \mathbb{R}^d$ that maps the $l + 1$ vertices of Δ_l to these $l + 1$ points. Thus the Tverberg theorem can be reformulated as follows:

Tverberg Theorem. *Let f be a linear map from the N -dimensional simplex Δ_N to $\mathbb{R}^d$. If $N = (d + 1)(r - 1)$ then there are r disjoint faces of Δ_N whose images have a common point.*

The following conjecture is a generalization of Tverberg Theorem to arbitrary continuous maps.

The topological Tverberg conjecture. *Let f be a continuous map from the N -dimensional simplex Δ_N to $\mathbb{R}^d$. If $N = (d + 1)(r - 1)$ then there are r disjoint faces of Δ_N whose images have a common point.*

The topological Tverberg conjecture was considered a central unsolved problem of topological combinatorics. For a prime number r the conjecture was proved by Bárány, Shlosman and Szűcs ([1]) and it was extended for a prime power r by Özaydin (unpublished) ([15]) and Volovikov ([19]). This result is known as the *topological Tverberg theorem*. Recently, in [6], Frick presents surprising counterexamples to the topological Tverberg conjecture for any r that is not a power of a prime and dimensions $d \geq 3r + 1$ (see also [3]). Although, the conjecture is not true for an integer $r \geq 6$ that is not a prime power, it is possible to prove a weak version of the topological Tverberg conjecture, more precisely, in this paper we show that if r is a natural number with prime factorization $r = p_1^{n_1} \cdots p_k^{n_k}$ then there is, for each $j = 1, \dots, k$, a set with r closed sides mutually disjoint of Δ_N which can be divided into $\frac{r}{p_j^{n_j}}$ subsets, each one having $p_j^{n_j}$ elements, whose images have a common point. Specifically, we prove the

following *Topological Tverberg type theorem for manifolds* and for any natural number r .

Theorem 5.1. *Let $d \geq 1$ a natural number. Consider a natural number r with prime factorization $r = p_1^{n_1} \cdots p_k^{n_k}$ and set $N = (r-1)(d+1)$. Let $f : \partial\Delta_N \rightarrow M$ be a continuous mapping into a compact d -dimensional topological manifold. Then, for each $j = 1, \dots, k$, among the sides of Δ_N there are $r = q_j r_j$, where $r_j = p_j^{n_j}$, and $q_j = \frac{r}{r_j}$, mutually disjoint closed sides $\sigma_{1_1}, \dots, \sigma_{1_{r_j}}; \dots; \sigma_{i_1}, \dots, \sigma_{i_{r_j}}; \dots; \sigma_{q_{j_1}}, \dots, \sigma_{q_{j_{r_j}}}$, such that*

$$f(\sigma_{i_1}) \cap \cdots \cap f(\sigma_{i_{r_j}}) \neq \emptyset, \text{ for each } i = 1, \dots, q_j.$$

Definition 5.2 (Index). *Let p be a prime. We suppose the p -torus $G = \mathbb{Z}_p^k = \mathbb{Z}_p \times \cdots \times \mathbb{Z}_p$ (k factors) acting freely on a paracompact space X . The covering $X \rightarrow X/G$ is induced from the universal covering $EG \rightarrow BG$ by means of a classifying map $c : X/G \rightarrow BG$, defined uniquely up to homotopy. We say that the **index** of X is greater than or equal to N (abbreviated by $\text{ind } X \geq N$) if $c^* : H^N(BG; \mathbb{Z}_p) \rightarrow H^N(X/G; \mathbb{Z}_p)$ is a monomorphism.*

Consider $G = \mathbb{Z}_{p_1}^{n_1} \times \cdots \times \mathbb{Z}_{p_k}^{n_k}$, where $\mathbb{Z}_{p_j}^{n_j} = \mathbb{Z}_{p_j} \times \cdots \times \mathbb{Z}_{p_j}$ (n_j factors), $j = 1, \dots, k$. We suppose G acts freely on a paracompact space X . Let $f : X \rightarrow M$ be a continuous mapping into a d -dimensional topological manifold.

Lemma 5.3. *Let M be a compact d -dimensional topological manifold orientable. Suppose the homomorphism $f^* : H^i(M; \mathbb{Z}_{p_j}) \rightarrow H^i(X; \mathbb{Z}_{p_j})$ is trivial for $i \geq 1$, and $\text{ind } X \geq N \geq d(r - q_j)$, for each $j = 1, \dots, k$, where $q_j = \frac{r}{p_j^{n_j}}$. Then*

$$\text{ind } A\left(f, \mathbb{Z}_{p_j}^{n_j}, G\right) \geq N - d(q - q_j).$$

Proof. We denote by $a_1, \dots, a_{q_j}$ a set of representatives of the left lateral classes of $G/\mathbb{Z}_{p_j}^{n_j}$. We define, for each $i = 1, \dots, q_j$, $g_i : X \rightarrow X$ by $g_i(x) = a_i x$. Consider the map $F : X \rightarrow M^{q_j}$ defined by

$$F = (f_1 \times \cdots \times f_{q_j}) \circ D,$$

where $(f_1 \times \cdots \times f_{q_j})(x_1, \dots, x_{q_j}) = (f_1(x_1), \dots, f_{q_j}(x_{q_j}))$, $D : X \rightarrow X^{q_j}$ is the diagonal map and $f_i = f \circ g_i$.

We have $F^* : H^i(M^{q_j}; \mathbb{Z}_{p_j}) \rightarrow H^i(X; \mathbb{Z}_{p_j})$ trivial for $i \geq 1$, therefore the index of $A(F) = \{x \in X : F(x) = F(gx) \forall g \in \mathbb{Z}_{p_j}^{n_j}\}$ is greater than or equal to $N - q_j d(p_j^{n_j} - 1)$ (see [18, Theorem 1]). Since $A(F) \subset A(f, \mathbb{Z}_{p_j}^{n_j}, G)$ and the inclusion $A(F) \hookrightarrow A(f, \mathbb{Z}_{p_j}^{n_j}, G)$ is an equivariant map we have $\text{ind } A(f, \mathbb{Z}_{p_j}^{n_j}, G) \geq A(F)$. Then

$$\text{ind } A\left(f, \mathbb{Z}_{p_j}^{n_j}, G\right) \geq N - d(r - q_j).$$

■

Proof of Theorem 5.1. We consider the CW -complex $Y_{N,r}$ that consists of points $(y_1, \dots, y_r)$, y_i in the boundary $\partial\Delta_N$ of the simplex Δ_N , that have mutually disjoint closed faces. It is known that for all natural numbers r and N , where $N \geq r + 1$, $Y_{N,r}$ is $(N - r)$ -connected (see [1]). Let $G = \{g_1, \dots, g_r\}$ be a fixed enumeration of elements of G , we define a G -action on $Y_{N,r} \subset (\Delta_N)^q$ as follows: for each $g_i \in G$ and for each $(y_1, \dots, y_r) \in Y_{N,r}$

$$g_i(y_1, \dots, y_r) = (y_{\phi_{g_i}(1)}, \dots, y_{\phi_{g_i}(r)}),$$

where the permutation ϕ_{g_i} , is defined by $\phi_{g_i}(k) = j$, $g_k g_i = g_j$. Then G acts freely on $Y_{N,r}$, since $Y_{N,r}$ consists of points $(y_1, \dots, y_r)$, $y_i \in \partial\Delta_N$ that have mutually disjoint closed faces.

Let $\tilde{f} : Y_{N,r} \rightarrow M$ given by $\tilde{f}(y_1, \dots, y_r) = f(y_1)$, that is, $\tilde{f} = f \circ \pi_1$ where $\pi_1 : Y_{N,r} \rightarrow \partial\Delta^N$ is the projection on the 1-th coordinate. Since $\tilde{f}^* : H^i(M; \mathbb{Z}_{p_j}) \rightarrow H^i(Y_{N,r}; \mathbb{Z}_{p_j})$ is trivial for $i \geq 1$ and $\text{ind } Y_{N,r} \geq d(r-1) > d(r-q_j)$ then, according to Lemma 5.3, the set $A(\tilde{f}, \mathbb{Z}_{p_j}^{n_j}, G)$ is not empty, for $j = 1, \dots, k$.

Let $H = \mathbb{Z}_{p_j}^{n_j} = \{h_1, \dots, h_{r_j}\}$ be a fixed enumeration of elements of $H = \mathbb{Z}_{p_j}^{n_j} \subset G$. We denote by $a_1, \dots, a_{q_j}$ a set of representatives of the left lateral classes of $G/\mathbb{Z}_{p_j}^{n_j}$ then, for each $i = 1, \dots, q_j$, $a_i h_1^{-1} = g_{l_1}, \dots, a_i h_{r_j}^{-1} = g_{l_{r_j}}$ are elements of G . Thus, if $y = (y_1, \dots, y_r) \in A(\tilde{f}, \mathbb{Z}_{p_j}^{n_j}, G)$ then

$$\tilde{f}(g_{l_1} \cdot (y_1, \dots, y_r)) = \dots = \tilde{f}(g_{l_{q_j}} \cdot (y_1, \dots, y_r))$$

that is

$$f(y_{\phi_{l_1}(1)}) = \dots = f(y_{\phi_{l_{q_j}}(1)}).$$

Therefore, for each $j = 1, \dots, k$, among the sides of Δ_N there are $r = q_j r_j$ mutually disjoint closed sides $\{\sigma_{i_1}, \dots, \sigma_{i_{r_j}}\}_{i=1}^{q_j}$, such that

$$f(\sigma_{i_1}) \cap \dots \cap f(\sigma_{i_{r_j}}) \neq \emptyset,$$

for each $i = 1, \dots, q_j$. ■

Let us observe that since the d -dimensional Euclidean space $\mathbb{R}^d$ is homeomorphic to the interior of the closed d -dimensional ball, Theorem 5.1 hold also for maps into $\mathbb{R}^d$, and we have the following *weak version of the topological Tverberg conjecture or topological Tverberg type theorem for any interger r* .

Theorem 5.4 (Topological Tverberg type theorem for any interger r). *Let $r \geq 2$, $d \geq 1$ be integers and $N = (r-1)(d+1)$. Consider $r = r_1 \dots r_k$ the prime factorization of r and denote $q_j = r/r_j$, $j = 1, \dots, k$. Then for any continuous map $f : \Delta_N \rightarrow \mathbb{R}^d$, for each $j = 1, \dots, k$, there are $r = q_j r_j$ pairwise disjoint faces $\{\sigma_{i_1}, \dots, \sigma_{i_{r_j}}\}_{i=1}^{q_j}$ such that*

$$f(\sigma_{i_1}) \cap \dots \cap f(\sigma_{i_{r_j}}) \neq \emptyset, \text{ for each } i = 1, \dots, q_j.$$

Remark 5.5. Let us note that if we consider r a prime power in Theorem 5.4, we obtain the topological Tverberg theorem for prime powers.

Now, by Theorem 5.4 and using similar method as in [3], we have the following *Generalized Van Kampen-Flores type theorem for any interger r or a weak version of the Generalized Van Kampen-Flores theorem.* In [3, Theorem 4.2], Blagojevic, Frick and Ziegler proved that the Generalized Van Kampen-Flores theorem does not hold in general.

Theorem 5.6 (Generalized Van Kampen-Flores type theorem for any r). *Let $d \geq 1$ a natural number. Consider a natural number r with prime factorization $r = r_1 \cdots r_k$, $r_1 < \cdots < r_k$, set $N = (r - 1)(d + 2)$ and let $l \geq \lceil \frac{r-1}{r_k} d \rceil$. Let $f : \Delta_N \rightarrow \mathbb{R}^d$ be a continuous mapping. Then, there are $r = q_k r_k$ pairwise disjoint faces $\{\sigma_{i_1}, \dots, \sigma_{i_{r_j}}\}_{i=1}^{q_k}$ of the l -th skeleton $\Delta_N^{(l)}$, such that*

$$f(\sigma_{i_1}) \cap \cdots \cap f(\sigma_{i_{r_j}}) \neq \emptyset, \text{ for each } i = 1, \dots, q_k.$$

Proof. Let $g : \Delta_N \rightarrow \mathbb{R}^{d+1}$ be a continuous function defined by $g(x) = (f(x), \text{dist}(x, \Delta_N^{(l)}))$. Then, we can apply Theorem 5.1 to function g which results in a collection of points

$$x_{1_1}, \dots, x_{1_{r_k}}; \dots; x_{i_1}, \dots, x_{i_{r_k}}; \dots; x_{q_{k_1}}, \dots, x_{q_{k_{r_k}}},$$

such that $\{x_{i_1}, \dots, x_{i_{r_k}}\}_{i=1}^{q_k}$ are points in the pairwise disjoint faces $\{\sigma_{i_1}, \dots, \sigma_{i_{r_j}}\}_{i=1}^{q_k}$ with $f(x_{i_1}) = \cdots = f(x_{i_{r_k}})$ and $\text{dist}(x_{i_1}, \Delta_N^{(l)}) = \cdots = \text{dist}(x_{i_{r_k}}, \Delta_N^{(l)})$, for each $i = 1, \dots, q_k$. We can suppose that the all σ_{i_s} 's are inclusion-minimal with the property that $x_{i_s} \in \sigma_{i_s}$, that is, σ_{i_s} is the unique face with x_{i_s} in its relative interior.

Now, for each $i = 1, \dots, q_k$ fixed, suppose that one of the faces $\sigma_{i_1}, \dots, \sigma_{i_{r_k}}$ is in $\Delta_N^{(l)}$, e.g. σ_{i_1} , then $\text{dist}(x_{i_1}, \Delta_N^{(l)}) = 0$, which imply that $\text{dist}(x_{i_1}, \Delta_N^{(l)}) = \cdots = \text{dist}(x_{i_{r_k}}, \Delta_N^{(l)}) = 0$, and consequently, all faces $\sigma_{i_1}, \dots, \sigma_{i_{r_k}}$ are in $\Delta_N^{(l)}$.

Let us suppose the contrary, that no σ_{i_s} is in $\Delta_N^{(l)}$, i.e., $\dim \sigma_{i_1} \geq l + 1, \dots, \dim \sigma_{i_{r_k}} \geq l + 1$. Since the faces $\sigma_{i_1}, \dots, \sigma_{i_{r_k}}$ are pairwise disjoint we have

$$\begin{aligned} N + 1 = |\Delta_N| &\geq |\sigma_{i_1}| + \cdots + |\sigma_{i_{r_k}}| \\ &\geq r_k(l + 2) \geq r_k \left(\left\lceil \frac{r-1}{r_k} d \right\rceil + 2 \right) \geq (r - 1)(d + 2) + 2 = N + 2, \end{aligned}$$

which is a contradiction and thus one of the faces $\sigma_{i_1}, \dots, \sigma_{i_{r_k}}$ is in $\Delta_N^{(l)}$ and consequently all faces $\sigma_{i_1}, \dots, \sigma_{i_{r_k}}$ are in $\Delta_N^{(l)}$. $\blacksquare$

Remark 5.7. Let us observe that if we consider r a prime power in Theorem 5.6, we obtain the Generalized Van Kampen-Flores theorem for prime powers proved by Blagojevic, Frick and Ziegler in [3, Theorem 3.2].

References

- [1] Bárány, I.; Shlosman, S. B.; Szűcz, A., *On a Topological generalization of a theorem of Tverberg*, J. London Math. Soc. **23** (1981), 158–164.
- [2] Birch, B. J., *On $3N$ points in a plane*, Math. Proc. Cambridge Phil. Soc. **55** (1959), 289–293.
- [3] Blagojevic, P., Frick, F., Ziegler, G. *Barycenters of Polytope Skeleta and Counterexamples to the Topological Tverberg Conjecture, via Constraints* eprint arXiv:1510.07984.
- [4] Borsuk, K., *Drei Sätze über die n -dimensionale euklidische Sphäre*, Fund. Math. **20** 177–190, 1933.
- [5] Conner, P.E.; Floyd E.E. *Differentiable Periodic Maps*. Ergebnisse der Math. Springer-Verlag Berlin **33**, 1964.
- [6] Frick, F., *Counterexamples to the topological Tverberg conjecture*, Preprint, 3 pages, arXiv:1502.00947, February 2015; to appear in Oberwolfach Reports, (1)11, 2015.
- [7] Gonçalves, D.L.; Pergher, Pedro L.Q., $\mathbb{Z}_p$ -coincidence for maps of spheres into CW-complexes, Kobe J. Math. **15**, (1998), 191–195.
- [8] Gonçalves, D.L.; Jaworowski, J.; Pergher, Pedro L.Q., *Measuring the size of the coincidence set*, Topology Appl. **125** (2002), 465–470.
- [9] Gonçalves, D.L.; Jaworowski, J.; Pergher, Pedro L.Q., A.Yu. Volovikov, *Coincidences for maps of spaces with finite group actions*. Topology Appl. **145** (2004), no. 1-3, 61–68.
- [10] Izydorek, M.; Jaworowski, J., *Antipodal coincidence for maps of spheres into complexes*, Proc. Amer. Math. Soc. **123** (6) (1995) 1947–1950.
- [11] De Mattos, Denise; Dos Santos, Edivaldo L., *On nonsymmetric theorems for (H, G) -coincidences*. Topological Methods in Nonlinear Analysis, v. 33, (2009) 105–120.
- [12] Milnor, J. W.; Stasheff, J. D., *Characteristic classes*, Princeton Univ. Press, Princeton, N.J. (1974).
- [13] Munkholm, H.J., *Borsuk-Ulam theorem for proper $\mathbb{Z}_p$ -actions on $(\text{mod } p \text{ homology})$ n -spheres*. Math. Scan. **24** (1969) 167–185.
- [14] Nakaoka, M., *Generalization of Borsuk-Ulam theorem*. Osaka J. Math. **7** (1970) 423–441.
- [15] Özaydin, M., *Equivariant maps for the symmetric group*, unpublished, 17 pages, availableonline at <http://minds.wisconsin.edu/handle/1793/63829> (1987).

- [16] Tverberg, H., *A generalization of Radon's Theorem*, J. London Math. Soc. 41 (1966), 123-128.
- [17] Volovikov, A. Yu., *Mappings of free $\mathbb{Z}_p$ -spaces into manifolds*, Izv. Akad. Nauk SSSR Ser. Math., (46) n.1, (1992) 36-55.
- [18] Volovikov, A. Yu., *A Bourgin-Yang-type theorem for $\mathbb{Z}_p^n$ -action*. (Russian) Mat. Sb. 183 (1992), no. 7, 115-144; translation in Russian Acad. Sci. Sb. Math. 76 (1993), no. 2, 361-387.
- [19] Volovikov, A. Yu., *On a topological generalization of Tverberg's theorem*, Mat. Zametki, (59) n.3 (1996), 454-456.

Denise de Mattos

E-mail address deniseml@icmc.usp.br

Universidade de São Paulo-USP-ICMC, Departamento de Matemática, Caixa Postal 668, 13560-970, São Carlos-SP, Brazil

Edivaldo L. dos Santos

E-mail address edivaldo@dm.ufscar.br

Departamento de Matemática, Universidade Federal de São Carlos, Centro de Ciências Exatas e Tecnologia, CP 676, CEP 13565-905, São Carlos - SP, Brazil.

Taciana O. Souza

E-mail address tacioli@ufu.br

Faculdade de Matemática, Universidade Federal de Uberlândia, Campus Santa Mônica - Bloco 1F - Sala 1F120, Av. João Naves de Ávila, 2121, Uberlândia, MG, CEP: 38.408-100, Brazil